arXiv:1108.3009v1 [math.FA] 15 Aug 2011

Operator inequalities dealing with operator equations *

Jian Shi[1] Zongsheng Gao[2]
[1, 2]LMIB & School of Mathematics and Systems Science,
Beihang University, Beijing, 100191, China

Abstract In this paper, we study the existence of solutions of some kinds of operator equations via operator inequalities. First, we investigate characterizations of operator order $A \geqslant B > 0$ and chaotic operator order $\log A \geqslant \log B$ for positive definite operators A, B in terms of operator equations, and generalize the results in [5]. Then, we introduce applications of complete form of Furuta inequality in operator equations. Some kinds of operator equations are researched and related characterizations of solutions are proved.

Keywords: Furuta inequality; Operator order ; Chaotic operator order; Operator equation; Complete form of Furuta inequality
Mathematics Subject Classification: 47A63.

1 Introduction

A bounded linear operator A on a Hilbert space H is said to be positive if $\langle Ax, x\rangle \geqslant 0$ for all $x \in H$. $A \geqslant 0$ and $A > 0$ mean a positive semidefinite operator and a positive definite operator, respectively. If $X > 0$, $\log X$ is defined as $\log X = lim_{a\to +0}\frac{X^a - I}{a}$. We call the relation $\log A \geqslant \log B$ chaotic operator order (in symbol: $A \gg B$).

In 1987 and 1995, Furuta showed the following operator inequalities:

Theorem 1.1 (Furuta Inequality) [2]. If $A \geqslant B \geqslant O$, then for each $r \geqslant 0$,

$$(B^{\frac{r}{2}}A^pB^{\frac{r}{2}})^{\frac{1}{q}} \geqslant (B^{\frac{r}{2}}B^pB^{\frac{r}{2}})^{\frac{1}{q}}, \tag{1.1}$$

$$(A^{\frac{r}{2}}A^pA^{\frac{r}{2}})^{\frac{1}{q}} \geqslant (A^{\frac{r}{2}}B^pA^{\frac{r}{2}})^{\frac{1}{q}} \tag{1.2}$$

hold for $p \geqslant 0$, $q \geqslant 1$ with $(1 + r)q \geqslant p + r$.

Theorem 1.2 (Grand Furuta Inequality) [3]. If $A \geqslant B \geqslant O$ with $A > O$, then for

---

*This work is supported by National Natural Science Fund of China (10771011).

[1] Corresponding author, E-mail addresses: shijian@ss.buaa.edu.cn (J. Shi).

[2] E-mail addresses: zshgao@buaa.edu.cn (Z. Gao).

$t \in [0,1]$ and $p \geqslant 1$,

$$A^{1-t+r} \geqslant (A^{\frac{r}{2}}(A^{-\frac{t}{2}}B^pA^{-\frac{t}{2}})^sA^{\frac{r}{2}})^{\frac{1-t+r}{(p-t)s+r}} \tag{1.3}$$

for $s \geqslant 1$ and $r \geqslant t$.

In 1996, Tanahashi proved the conditions for p and q in Furuta Inequality are the best possible if $r \geqslant 0$ in [7]; In 2000, he proved the outer exponent value of Grand Furuta Inequality is the best possible in [8]. In 2008, Yuan et al. proved a new operator inequality, which is called Complete form of Furuta Inequality:

Theorem 1.3 (Complete form of Furuta Inequality) [11]. Let $A \geqslant B \geqslant O$, $r \geqslant 0$, $p > p_0 \geqslant 0$, and $s = min\{p, 2p_0 + min\{1, r\}\}$. Then

$$(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{\frac{s+r}{p_0+r}} \geqslant (A^{\frac{r}{2}}B^pA^{\frac{r}{2}})^{\frac{s+r}{p+r}}. \tag{1.4}$$

Yuan et al. also proved the optimality of the outer exponent of Complete form of Furuta Inequality under the condition of $\{2p_0 + min\{1, r\}\} \geqslant p$ or $r \geqslant 1$ with $2p_0 + 1 < p$ in [11]. There were many related studies on Complete form of Furuta Inequality, such as [12], [13].

Recently, some kinds of operator equations have been shown and given deep discussion via Furuta inequality and grand Furuta inequality (see [5], [9]). In [5], C.-S. Lin proved the following two theorems, which the characterizations of operator order $A \geqslant B > 0$ and chaotic operator order $A \gg B$ for positive definite operator A, B in terms of operator equations are discussed, respectively:

Theorem 1.4 [5]. For $r \geqslant 0$ and a nonnegative integer $n \geqslant 0$ such that $(1 + r)(n + 1) = p + r$ (so, $p \geqslant 1$), the following assertions are equivalent.

(A1) $A \geqslant B > O$;

(A2) $A^{1+r} \geqslant (A^{r/2}B^pA^{r/2})^{\frac{1}{n+1}}$;

(A3) $(B^{r/2}A^pB^{r/2})^{\frac{1}{n+1}} \geqslant B^{1+r}$;

(A4) there exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$B^p = A^{1/2}S(A^{1+r}S)^nA^{1/2} = A^{1/2}(SA^{1+r})^nSA^{1/2}$;

(A5) there exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$A^p = B^{1/2}S^{-1}(B^{1+r}S^{-1})^nB^{1/2} = B^{1/2}(S^{-1}B^{1+r})^nS^{-1}B^{1/2}$.

Theorem 1.5 [5]. For $p, r > 0$ and any integer $n \geqslant 1$ such that $r(n + 1) = p + r$, the following assertions are equivalent.

(B1) $A \gg B$;

(B2) $A^r \geqslant (A^{r/2}B^pA^{r/2})^{\frac{1}{n+1}}$;

(B3) $(B^{r/2}A^pB^{r/2})^{\frac{1}{n+1}} \geqslant B^r$;

(B4) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$B^p = T(A^rT)^n = (TA^r)^nT$;

(B5) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$A^p = T^{-1}(B^rT^{-1})^n = (T^{-1}B^r)^nT^{-1}$.

In this paper, we will show generalized characterizations in terms of operator equations due to Theorem 1.4 and Theorem 1.5. Then, we will give some applications of Complete form of Furuta Inequality in operator equations.

In order to prove our results, let's recall some well-known theorems:

Theorem 1.7 (Löwner-Heinz inequality) [6], [4]. $A \geqslant B \geqslant 0$ ensures $A^\alpha \geqslant B^\alpha$ for any $\alpha \in [0, 1]$.

Theorem 1.7 assures that $A \geqslant B > 0 \Rightarrow A \gg B$.

Theorem 1.8 (Douglas's majorization and factorization theorem) [1]. The following assertions are equivalent for any operators A and B.

(a) range(A)$\subseteq$ range (B);

(b) $AA^* \leqslant \lambda BB^*$ for some $\lambda \geqslant 0$ (majorization);

(c) There exists a bounded operator C such that $A = BC$ (factorization).

Moreover, if (a), (b) and (c) are valid, then there exists a unique operator C such that

(i)$\|C\|^2 = inf\{\mu | AA^* \leqslant BB^*\}$;

(ii) Null(A)=Null(C);

(iii) range(C)$\subseteq$ range(B^*).

Theorem 1.9 [10]. For $A, B > O$, $A \geqslant B$ if and only if

$$A^{1+t+r} \geqslant (A^{r/2}(A^{t/2}B^pA^{t/2})^sA^{r/2})^{(1+t+r)/((p+t)s+r)}$$

holds for $p \geqslant 1$, $t \geqslant 0$, $r \geqslant 0$ and $s \geqslant \frac{1+t}{p+t}$.

Theorem 1.10 [10]. For $A, B > O$, log $A \geqslant$ log B if and only if

$$A^{t+r} \geqslant (A^{r/2}(A^{t/2}B^pA^{t/2})^sA^{r/2})^{(t+r)/((p+t)s+r)}$$

holds for $p > 0$, $t \geqslant 0$, $r \geqslant 0$ and $s \geqslant \frac{t}{p+t}$.

2 The characterizations of operator order $A \geqslant B > O$ and chaotic operator order $A \gg B$

In this section , we will show further development of characterizations of operator order $A \geqslant B > 0$ and chaotic operator order $\log A \geqslant \log B$ for positive definite operators A, B in terms of operator equations due to Lin's results in [5].

Throughout this section we assume that A, B are positive definite operators.

Theorem 2.1. For $r \geqslant 0$, $t \geqslant 0$, $p \geqslant 1$ and a nonnegative n such that $(p+t)s + r = (n+1)\cdot(1+t+r)$ (so, $s \geqslant \frac{1+t}{p+t}$), the following assertions are equivalent.

(C1) $A \geqslant B > O$;

(C2) $A^{1+t+r} \geqslant (A^{r/2}(A^{t/2}B^pA^{t/2})^sA^{r/2})^{\frac{1}{n+1}}$;

(C3) $(B^{r/2}(B^{t/2}A^pB^{t/2})^sB^{r/2})^{\frac{1}{n+1}} \geqslant B^{1+t+r}$;

(C4) there exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$$\begin{aligned} B^p &= A^{-\frac{t}{2}}(A^{\frac{1+t}{2}}(SA^{1+t+r})^nSA^{\frac{1+t}{2}})^{\frac{1}{s}}A^{-\frac{t}{2}} \\ &= A^{-\frac{t}{2}}(A^{\frac{1+t}{2}}S(A^{1+t+r}S)^nA^{\frac{1+t}{2}})^{\frac{1}{s}}A^{-\frac{t}{2}} \ ; \end{aligned}$$

(C5) there exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$$\begin{aligned} A^p &= B^{-\frac{t}{2}}(B^{\frac{1+t}{2}}(S^{-1}B^{1+t+r})^nS^{-1}B^{\frac{1+t}{2}})^{\frac{1}{s}}B^{-\frac{t}{2}} \\ &= B^{-\frac{t}{2}}(B^{\frac{1+t}{2}}S^{-1}(B^{1+t+r}S^{-1})^nB^{\frac{1+t}{2}})^{\frac{1}{s}}B^{-\frac{t}{2}}. \end{aligned}$$

Proof.

(C1) $\Rightarrow$ (C2) is obvious by Theorem 1.9.

(C2) $\Rightarrow$ (C1). Take $n = 0$, $t = 0$, $r = 0$ in $(p+t)s + r = (n+1)\cdot(1+t+r)$ and (C2), then $ps = 1$ and $A \geqslant B$.

(C2) $\Leftrightarrow$ (C3) holds by $A \geqslant B > O \Leftrightarrow B^{-1} \geqslant A^{-1} > O$.

(C2) $\Rightarrow$ (C4). According to Theorem 1.8, there exists C with $\|C\| \leqslant 1$ such that $(A^{r/2}(A^{t/2}B^pA^{t/2})^sA^{r/2})^{\frac{1}{2(n+1)}} = A^{\frac{1+t+r}{2}}C = C^*A^{\frac{1+t+r}{2}}$.

Take $S = CC^*$, by $\|C\| \leqslant 1$ and C is determined by A and B, we have $\|S\| \leqslant 1$, S is unique, and $(A^{r/2}(A^{t/2}B^pA^{t/2})^sA^{r/2})^{\frac{1}{n+1}} = A^{\frac{1+t+r}{2}}SA^{\frac{1+t+r}{2}}$. Therefore, the following equalities hold.

$$A^{r/2}(A^{t/2}B^pA^{t/2})^sA^{r/2} = (A^{\frac{1+t+r}{2}}SA^{\frac{1+t+r}{2}})^{n+1} = A^{\frac{1+t+r}{2}}(SA^{1+t+r})^nSA^{\frac{1+t+r}{2}}; \quad (2.1)$$

$$A^{r/2}(A^{t/2}B^pA^{t/2})^sA^{r/2} = (A^{\frac{1+t+r}{2}}SA^{\frac{1+t+r}{2}})^{n+1} = A^{\frac{1+t+r}{2}}S(A^{1+t+r}S)^nA^{\frac{1+t+r}{2}}. \quad (2.2)$$

By (2.1) and (2.2), $(A^{t/2}B^pA^{t/2})^s = A^{\frac{1+t}{2}}(SA^{1+t+r})^nSA^{\frac{1+t}{2}} = A^{\frac{1+t}{2}}S(A^{1+t+r}S)^nA^{\frac{1+t}{2}}$, then (C4) is obtained.

(C4) $\Rightarrow$ (C2).

$$\begin{aligned}
& (A^{r/2}(A^{t/2}B^pA^{t/2})^sA^{r/2})^{\frac{1}{n+1}} \\
= \ & (A^{r/2}(A^{t/2}A^{-t/2}(A^{\frac{1+t}{2}}(SA^{1+t+r})^nSA^{\frac{1+t}{2}})^{1/s}A^{-t/2}A^{t/2})^sA^{r/2})^{\frac{1}{n+1}} \\
= \ & (A^{\frac{1+t+r}{2}}(SA^{1+t+r})^nSA^{\frac{1+t+r}{2}})^{\frac{1}{n+1}} \\
= \ & ((A^{\frac{1+t+r}{2}}SA^{\frac{1+t+r}{2}})^{n+1})^{\frac{1}{n+1}} \\
= \ & A^{\frac{1+t+r}{2}}SA^{\frac{1+t+r}{2}} \\
\leqslant \ & A^{1+t+r}.
\end{aligned}$$

The first equality is due to the equation of (C4), and the last inequality is due to $O < S \leqslant \|S\|I \leqslant I$.

(C3) $\Rightarrow$ (C5). By (C3), we have $B^{-(1+t+r)} \geqslant (B^{-r/2}(B^{-t/2}A^{-p}B^{-t/2})^sB^{-r/2})^{\frac{1}{n+1}}$. According to Theorem 1.8, there exists C, such that $\|C\| \leqslant 1$, and $(B^{-r/2}(B^{-t/2}A^{-p}B^{-t/2})^sB^{-r/2})^{\frac{1}{2(n+1)}} = B^{-(1+t+r)/2}C = C^*B^{-(1+t+r)/2}$. Take $S = CC^*$, because $\|C\| \leqslant 1$ and C is uniquely determined by A and B, we have $\|S\| \leqslant 1$ and S is unique, and $(B^{-r/2}(B^{-t/2}A^{-p}B^{-t/2})^sB^{-r/2})^{\frac{1}{n+1}} = B^{-(1+t+r)/2}SB^{-(1+t+r)/2}$. Therefore, the following equality holds.

$$\begin{aligned}
& B^{r/2}(B^{t/2}A^pB^{t/2})^sB^{r/2} \\
= \ & (B^{-r/2}(B^{-t/2}A^{-p}B^{-t/2})^sB^{-r/2})^{-1} \\
= \ & (B^{-(1+t+r)/2}SB^{-(1+t+r)/2})^{-(n+1)} \\
= \ & (B^{(1+t+r)/2}S^{-1}B^{(1+t+r)/2})^{n+1} \\
= \ & B^{(1+t+r)/2}S^{-1}(B^{1+t+r}S^{-1})^nB^{(1+t+r)/2} \\
= \ & B^{(1+t+r)/2}(S^{-1}B^{1+t+r})^nS^{-1}B^{(1+t+r)/2}.
\end{aligned}$$

Then (C5) is obtained immediately.

(C5) ⇒ (C3).

$$\begin{aligned}
&(B^{r/2}(B^{t/2}A^pB^{t/2})^sB^{r/2})^{\frac{1}{n+1}} \\
=\ &(B^{r/2}(B^{t/2}B^{-t/2}(B^{\frac{1+t}{2}}S^{-1}(B^{1+t+r}S^{-1})^nB^{\frac{1+t}{2}})^{1/s}B^{-t/2}B^{t/2})^sB^{r/2})^{\frac{1}{n+1}} \\
=\ &(B^{\frac{1+t+r}{2}}S^{-1}(B^{1+t+r}S^{-1})^nB^{\frac{1+t+r}{2}})^{\frac{1}{n+1}} \\
=\ &((B^{\frac{1+t+r}{2}}S^{-1}B^{\frac{1+t+r}{2}})^{n+1})^{\frac{1}{n+1}} \\
=\ &B^{\frac{1+t+r}{2}}S^{-1}B^{\frac{1+t+r}{2}} \\
\geqslant\ &B^{1+t+r}.
\end{aligned}$$

The first equality is due to (C5), the last inequality is due to $O < S \leqslant \|S\|I \leqslant I \Rightarrow S^{-1} \geqslant I$.

Remark 2.1. The special case $r = 0$ and $s = 1$ of Theorem 2.1 just is Theorem 1.4. (A2), (A3), (A4), (A5) in Theorem 1.4 are extended to (C2), (C3), (C4), (C5) in Theorem 2.1, respectively.

Theorem 2.2. For p, $r > 0$, $t \geqslant 0$ and any integer $n \geqslant 1$ such that $(p+t)s + r = (n+1)\cdot(t+r)$, the following assertions are equivalent.

(D1) $A \gg B$;

(D2) $A^{t+r} \geqslant (A^{r/2}(A^{t/2}B^pA^{t/2})^sA^{r/2})^{\frac{1}{n+1}}$;

(D3) $(B^{r/2}(B^{t/2}A^pB^{t/2})^sB^{r/2})^{\frac{1}{n+1}} \geqslant B^{t+r}$;

(D4) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$\begin{aligned}
B^p &= A^{-\frac{t}{2}}(A^{\frac{t}{2}}(TA^{t+r})^nTA^{\frac{t}{2}})^{\frac{1}{s}}A^{-\frac{t}{2}} \\
&= A^{-\frac{t}{2}}(A^{\frac{t}{2}}T(A^{t+r}T)^nA^{\frac{t}{2}})^{\frac{1}{s}}A^{-\frac{t}{2}} \ ;
\end{aligned}$$

(D5) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$\begin{aligned}
A^p &= B^{-\frac{t}{2}}(B^{\frac{t}{2}}(T^{-1}B^{t+r})^nT^{-1}B^{\frac{t}{2}})^{\frac{1}{s}}B^{-\frac{t}{2}} \\
&= B^{-\frac{t}{2}}(B^{\frac{t}{2}}T^{-1}(B^{t+r}T^{-1})^nB^{\frac{t}{2}})^{\frac{1}{s}}B^{-\frac{t}{2}}.
\end{aligned}$$

Proof.

(D1) ⇒ (D2) is obvious by Theorem 1.10.

(D2) ⇒ (D1). Take $t = 0$, $r = \frac{1}{n}$, then $ps = 1$ and $A^{\frac{1}{n}} \geqslant (A^{\frac{1}{2n}}BA^{\frac{1}{2n}})^{\frac{1}{n+1}}$. Take logarithm of both sides and refining, we have $\log A \geqslant \frac{n}{n+1} \log\,(A^{\frac{1}{2n}}BA^{\frac{1}{2n}})$, and let $n \to \infty$, $\log A \geqslant \log B$ is obtained.

(D2) ⇔ (D3) is due to the fact that $A \gg B \Leftrightarrow B^{-1} \gg A^{-1}$.

The proofs of (D2) $\Leftrightarrow$ (D4) and (D3) $\Leftrightarrow$ (D5) are similar to the proofs of (C2) $\Leftrightarrow$ (C4) and (C3) $\Leftrightarrow$ (C5), and we omit them here.

Remark 2.2. The special case $t = 0$ and $s = 1$ of Theorem 2.2 just is Theorem 1.5. (B2), (B3), (B4), (B5) in Theorem 1.5 are extended to (D2), (D3), (D4), (D5) in Theorem 2.2, respectively.

Corollary 2.1. For $p > 0$, $t \geqslant 0$, the following assertions are equivalent.

(E1) $A \gg B$;

(E2) $A^{p+t} \geqslant (A^{p/2}(A^{t/2}B^pA^{t/2})^{(p+2t)/(p+t)}A^{p/2})^{1/2}$;

(E3) $(B^{p/2}(B^{t/2}A^pB^{t/2})^{(p+2t)/(p+t)}B^{p/2})^{1/2} \geqslant B^{p+t}$;

(E4) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$B^p = A^{-t/2}(A^{t/2}TA^{p+t}TA^{t/2})^{(p+t)/(p+2t)}A^{-t/2} \;;$$

(E5) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$A^p = B^{-t/2}(B^{t/2}T^{-1}B^{p+t}T^{-1}B^{t/2})^{(p+t)/(p+2t)}B^{-t/2}.$$

Proof. Take $n = 1$, $r = p$ in Theorem 2.2, then $s = (p + 2t)/(p + t)$.

Remark 2.3. If $t = 0$, Corollary 2.1 just is Corollary 3.1 in [5].

Corollary 2.2. For any positive integer m, n, the following assertions are equivalent.

(F1) $A \gg B$;

(F2) $A^{\frac{1}{m}+\frac{1}{n}} \geqslant (A^{\frac{1}{2n}}(A^{\frac{1}{2m}}BA^{\frac{1}{2m}})^{\frac{m+n+1}{m+1}}A^{\frac{1}{2n}})^{\frac{1}{n+1}}$;

(F3) $(B^{\frac{1}{2n}}(B^{\frac{1}{2m}}AB^{\frac{1}{2m}})^{\frac{m+n+1}{m+1}}B^{\frac{1}{2n}})^{\frac{1}{n+1}} \geqslant B^{\frac{1}{m}+\frac{1}{n}}$;

(F4) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$\begin{aligned} B &= A^{-\frac{1}{2m}}(A^{\frac{1}{2m}}(TA^{\frac{1}{m}+\frac{1}{n}})^nTA^{\frac{1}{2m}})^{\frac{m+1}{m+n+1}}A^{-\frac{1}{2m}} \\ &=A^{-\frac{1}{2m}}(A^{\frac{1}{2m}}T(A^{\frac{1}{m}+\frac{1}{n}}T)^nA^{\frac{1}{2m}})^{\frac{m+1}{m+n+1}}A^{-\frac{1}{2m}}; \end{aligned}$$

(F5) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$\begin{aligned} A &= B^{-\frac{1}{2m}}(B^{\frac{1}{2m}}(T^{-1}B^{\frac{1}{m}+\frac{1}{n}})^nT^{-1}B^{\frac{1}{2m}})^{\frac{m+1}{m+n+1}}B^{-\frac{1}{2m}} \\ &= B^{-\frac{1}{2m}}(B^{\frac{1}{2m}}T^{-1}(B^{\frac{1}{m}+\frac{1}{n}}T^{-1})^nB^{\frac{1}{2m}})^{\frac{m+1}{m+n+1}}B^{-\frac{1}{2m}}. \end{aligned}$$

Proof. Take $p = 1$, $r = \frac{1}{n}$, $t = \frac{1}{m}$ in Theorem 2.2, then $s = \frac{m+n+1}{m+1}$.

Remark 2.4. Let $m \to \infty$, Corollary 2.2 just is Corollary 3.2 in [5].

Corollary 2.3. For any positive integer m, n and some $a > 0$, the following assertions are equivalent.

(G1) $a^{n+1}A \gg B$;

(G2) $a^{\frac{m+n+1}{m+1}}A^{\frac{1}{m}+\frac{1}{n}} \geqslant (A^{\frac{1}{2n}}(A^{\frac{1}{2m}}BA^{\frac{1}{2m}})^{\frac{m+n+1}{m+1}}A^{\frac{1}{2n}})^{\frac{1}{n+1}}$;

(G3) $a^{\frac{m+n+1}{m+1}}(B^{\frac{1}{2n}}(B^{\frac{1}{2m}}AB^{\frac{1}{2m}})^{\frac{m+n+1}{m+1}}B^{\frac{1}{2n}})^{\frac{1}{n+1}} \geqslant B^{\frac{1}{m}+\frac{1}{n}}$;

(G4) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$B = a^{n+1}A^{-\frac{1}{2m}}(A^{\frac{1}{2m}}(TA^{\frac{1}{m}+\frac{1}{n}})^n TA^{\frac{1}{2m}})^{\frac{m+1}{m+n+1}}A^{-\frac{1}{2m}}$$
$$= a^{n+1}A^{-\frac{1}{2m}}(A^{\frac{1}{2m}}T(A^{\frac{1}{m}+\frac{1}{n}}T)^n A^{\frac{1}{2m}})^{\frac{m+1}{m+n+1}}A^{-\frac{1}{2m}} \;;$$

(G5) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$a^{n+1}A = B^{-\frac{1}{2m}}(B^{\frac{1}{2m}}(T^{-1}B^{\frac{1}{m}+\frac{1}{n}})^n T^{-1}B^{\frac{1}{2m}})^{\frac{m+1}{m+n+1}}B^{-\frac{1}{2m}}$$
$$= B^{-\frac{1}{2m}}(B^{\frac{1}{2m}}T^{-1}(B^{\frac{1}{m}+\frac{1}{n}}T^{-1})^n B^{\frac{1}{2m}})^{\frac{m+1}{m+n+1}}B^{-\frac{1}{2m}}.$$

Proof. Replace A by $a^{n+1}A$ in Corollary 2.2.

Remark 2.5. Let $m \to \infty$, Corollary 2.3 just is Corollary 3.3 in [5].

Corollary 2.4. For any positive integer m and some $a > 0$, the following assertions are equivalent.

(H1) $a^2A \gg B$;

(H2) $a^{\frac{m+2}{m+1}}A^{1+\frac{1}{m}} \geqslant (A^{\frac{1}{2}}(A^{\frac{1}{2m}}BA^{\frac{1}{2m}})^{\frac{m+2}{m+1}}A^{\frac{1}{2}})^{\frac{1}{2}}$;

(H3) $a^{\frac{m+2}{m+1}}(B^{\frac{1}{2}}(B^{\frac{1}{2m}}AB^{\frac{1}{2m}})^{\frac{m+2}{m+1}}B^{\frac{1}{2}})^{\frac{1}{2}} \geqslant B^{1+\frac{1}{m}}$;

(H4) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$B = a^2A^{-\frac{1}{2m}}(A^{\frac{1}{2m}}TA^{1+\frac{1}{m}}TA^{\frac{1}{2m}})^{\frac{m+1}{m+2}}A^{-\frac{1}{2m}} \;;$$

(H5) there exists a unique $T > O$ with $\|T\| \leqslant 1$ such that

$$a^2A = B^{-\frac{1}{2m}}(B^{\frac{1}{2m}}T^{-1}B^{1+\frac{1}{m}}T^{-1}B^{\frac{1}{2m}})^{\frac{m+1}{m+2}}B^{-\frac{1}{2m}} \;.$$

Proof. Take $n = 1$ in Corollary 2.3.

Remark 2.6. Let $m \to \infty$, Corollary 2.4 just is Corollary 3.4 in [5].

3 Applications of Complete form of Furuta Inequality in Operator Equations

In this section, we will introduce some applications of Complete form of Furuta Inequality in operator equations. Several kinds of operator equations will be researched and related characterizations of solutions will be proved.

Throughout this section we assume A and B are positive definite operators.

Theorem 3.1. If $1 \geqslant r \geqslant 0$, $p > p_0 \geqslant 0$, for a nonnegative integer n such that $p + r = (n+1)((2p_0 + r) + r)$, the following assertions are equivalent.

(3-1) $A \geqslant B > O$;

(3-2) $(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^2 \geqslant (A^{\frac{r}{2}}B^{p}A^{\frac{r}{2}})^{\frac{1}{n+1}}$;

(3-3) There exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$$B^p = B^{p_0}A^{\frac{r}{2}}S(A^{\frac{r}{2}}B^{p_0}A^rB^{p_0}A^{\frac{r}{2}}S)^nA^{\frac{r}{2}}B^{p_0} = B^{p_0}A^{\frac{r}{2}}(SA^{\frac{r}{2}}B^{p_0}A^rB^{p_0}A^{\frac{r}{2}})^nSA^{\frac{r}{2}}B^{p_0}.$$

Proof. (3-1) $\Rightarrow$ (3-2) is obvious by Theorem 1.3.

(3-2) $\Rightarrow$ (3-1). Take $p_0 = 0$, $r = 1$, $n = 0$ in $p + r = (n+1)((2p_0 + r + r)$ and (3-2), then $p = 1$ and $A^2 \geqslant A^{\frac{1}{2}}BA^{\frac{1}{2}}$.

(3-2) $\Rightarrow$ (3-3). According to Theorem 1.8, there exists an operator C with $\|C\| \leqslant 1$ such that $(A^{\frac{r}{2}}B^pA^{\frac{r}{2}})^{\frac{1}{2(n+1)}} = (A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})C = C^*(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})$. Take $S = CC^*$, then $\|S\| = \|CC^*\| \leqslant 1$ and

$$(A^{\frac{r}{2}}B^pA^{\frac{r}{2}})^{\frac{1}{n+1}} = (A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})S(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}}). \tag{3.1}$$

By (3.1), S is unique and the following equality holds,

$$\begin{aligned} & A^{\frac{r}{2}}B^pA^{\frac{r}{2}} \\ = \ & ((A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})S(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}}))^{n+1} \\ = \ & A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}}S((A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^2S)^nA^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}} \\ = \ & A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}}(S(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^2)^nSA^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}}. \end{aligned}$$

By above equalities, we can obtain (3.3).

(3-3) $\Rightarrow$ (3-2).

$$\begin{aligned} & (A^{\frac{r}{2}}B^pA^{\frac{r}{2}})^{\frac{1}{n+1}} \\ = \ & (A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}}S(A^{\frac{r}{2}}B^{p_0}A^rB^{p_0}A^{\frac{r}{2}}S)^nA^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{\frac{1}{n+1}} \\ = \ & ((A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}}SA^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{n+1})^{\frac{1}{n+1}} \\ = \ & (A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})S(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}}) \\ \leqslant \ & (A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^2. \end{aligned}$$

The first equality is due to (3-3), and the inequality is due to $O < S \leqslant \|S\|I \leqslant I$.

Corollary 3.1. If $1 \geqslant r \geqslant 0$, $p > p_0 \geqslant 0$, for a nonnegative integer $n \geqslant 0$ such that $p + r = (n+1)((2p_0 + r) + r)$, the following assertions are equivalent.

(3-1) $A \geqslant B > O$;

(3-4) $(B^{\frac{r}{2}}A^pB^{\frac{r}{2}})^{\frac{1}{n+1}} \geqslant (B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^2$;

(3-5) There exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$A^p = A^{p_0}B^{\frac{r}{2}}S^{-1}(B^{\frac{r}{2}}A^{p_0}B^rA^{p_0}B^{\frac{r}{2}}S^{-1})^nB^{\frac{r}{2}}A^{p_0} = A^{p_0}B^{\frac{r}{2}}(S^{-1}B^{\frac{r}{2}}A^{p_0}B^rA^{p_0}B^{\frac{r}{2}})^nS^{-1}B^{\frac{r}{2}}A^{p_0}$.

Proof. Replace A and B by B^{-1} and A^{-1} , respectively, in (3-2) and (3-3), then (3-4) and (3-5) are obtained.

Theorem 3.2. If $r \geqslant 1$, $p > p_0 \geqslant 0$, for a nonnegative integer n such that $p + r = (n + 1)((2p_0 + 1) + r)$, the following assertions are equivalent.

(3-1) $A \geqslant B > O$;

(3-6) $(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{\frac{2p_0+1+r}{p_0+r}} \geqslant (A^{\frac{r}{2}}B^pA^{\frac{r}{2}})^{\frac{1}{n+1}}$;

(3-7) There exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$$\begin{aligned} B^p &= A^{-\frac{r}{2}}(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{\frac{2p_0+1+r}{2p_0+2r}}S((A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{\frac{2p_0+1+r}{p_0+r}}S)^n(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{\frac{2p_0+1+r}{2p_0+2r}}A^{-\frac{r}{2}} \\ &= A^{-\frac{r}{2}}(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{\frac{2p_0+1+r}{2p_0+2r}}(S(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{\frac{2p_0+1+r}{p_0+r}})^nS(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{\frac{2p_0+1+r}{2p_0+2r}}A^{-\frac{r}{2}}. \end{aligned}$$

Proof. (3-1) $\Rightarrow$ (3-6) is obvious by Theorem 1.3.

(3-6) $\Rightarrow$ (3-1). Take $n = 0$, $p_0 = 0$, $r = 1$ in $p + r = (n + 1)((2p_0 + 1) + r)$ and (3-6), then $p = 1$ and $A^2 \geqslant A^{\frac{1}{2}}BA^{\frac{1}{2}}$. (3-1) is obtained.

The proof of (3-6) $\Leftrightarrow$ (3-7) is similar to the proof of (3-2) $\Leftrightarrow$ (3-3) in Theorem 3.1, so we omit it here.

Corollary 3.2. If $r \geqslant 1$, $p > p_0 \geqslant 0$, for a nonnegative integer n such that $p + r = (n + 1)((2p_0 + 1) + r)$, the following assertions are equivalent.

(3-1) $A \geqslant B > O$;

(3-8) $(B^{\frac{r}{2}}A^pB^{\frac{r}{2}})^{\frac{1}{n+1}} \geqslant (B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^{\frac{2p_0+1+r}{p_0+r}}$;

(3-9) There exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$$\begin{aligned} A^p &= B^{-\frac{r}{2}}(B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^{\frac{2p_0+1+r}{2p_0+2r}}S^{-1}((B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^{\frac{2p_0+1+r}{p_0+r}}S^{-1})^n(B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^{\frac{2p_0+1+r}{2p_0+2r}}B^{-\frac{r}{2}} \\ &= B^{-\frac{r}{2}}(B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^{\frac{2p_0+1+r}{2p_0+2r}}(S^{-1}(B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^{\frac{2p_0+1+r}{p_0+r}})^nS^{-1}(B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^{\frac{2p_0+1+r}{2p_0+2r}}B^{-\frac{r}{2}}. \end{aligned}$$

Proof. Replaced A and B by B^{-1} and A^{-1} in (3-6) and (3-7) of Theorem 3.2, respectively, then (3-8) and (3-9) are obtained.

Theorem 3.3. If $r \geqslant 0$, $2p_0 + min\{1, r\} \geqslant p > p_0 \geqslant 0$, for a positive integer n such that $n(p+r) = (n+1)(p_0+r)$, the following assertions are equivalent.

(3-1) $A \geqslant B > O$;

(3-10) $(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{1+\frac{1}{n}} \geqslant A^{\frac{r}{2}}B^{p}A^{\frac{r}{2}}$;

(3-11) There exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$$(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{n+1} = ((A^{\frac{r}{2}}B^{p}A^{\frac{r}{2}})^{\frac{1}{2}}S^{-1}(A^{\frac{r}{2}}B^{p}A^{\frac{r}{2}})^{\frac{1}{2}})^{n} .$$

Proof. (3-1) ⇒ (3-10) is obvious by Theorem 1.3.

(3-10) ⇒ (3-1). Take $n = 1$, $p_0 = 0$, $r = 1$ in $n(p+r) = (n+1)(p_0+r)$ and (3-10), then $p = 1$ and $A^2 \geqslant A^{\frac{1}{2}}BA^{\frac{1}{2}}$. $A \geqslant B$ is obtained.

(3-10) ⇒ (3-11). By (3-10) we have $A^{-\frac{r}{2}}B^{-p}A^{-\frac{r}{2}} \geqslant (A^{-\frac{r}{2}}B^{-p_0}A^{-\frac{r}{2}})^{\frac{n+1}{n}}$. According to Theorem 1.8, there exists an operator C with $\|C\| \leqslant 1$, such that

$$(A^{-\frac{r}{2}}B^{-p_0}A^{-\frac{r}{2}})^{\frac{n+1}{2n}} = (A^{-\frac{r}{2}}B^{-p}A^{-\frac{r}{2}})^{\frac{1}{2}}C = C^*(A^{-\frac{r}{2}}B^{-p}A^{-\frac{r}{2}})^{\frac{1}{2}}.$$

Take $S = CC^*$, then $S > O$, $\|S\| \leqslant 1$ and the following equality,

$$(A^{-\frac{r}{2}}B^{-p_0}A^{-\frac{r}{2}})^{\frac{n+1}{n}} = (A^{-\frac{r}{2}}B^{-p}A^{-\frac{r}{2}})^{\frac{1}{2}}S(A^{-\frac{r}{2}}B^{-p}A^{-\frac{r}{2}})^{\frac{1}{2}}. \tag{3.2}$$

By (3.2), S is unique and $(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{n+1} = ((A^{\frac{r}{2}}B^{p}A^{\frac{r}{2}})^{\frac{1}{2}}S^{-1}(A^{\frac{r}{2}}B^{p}A^{\frac{r}{2}})^{\frac{1}{2}})^{n}$.

(3-11) ⇒ (3-10). Because of (3-11) and the fact that $S > O$ with $\|S\| \leqslant 1 \Rightarrow S^{-1} \geqslant I$, we have $(A^{\frac{r}{2}}B^{p_0}A^{\frac{r}{2}})^{1+\frac{1}{n}} = (A^{\frac{r}{2}}B^{p}A^{\frac{r}{2}})^{\frac{1}{2}}S^{-1}(A^{\frac{r}{2}}B^{p}A^{\frac{r}{2}})^{\frac{1}{2}} \geqslant A^{\frac{r}{2}}B^{p}A^{\frac{r}{2}}$.

Corollary 3.3. If $r \geqslant 0$, $2p_0 + min\{1, r\} \geqslant p > p_0 \geqslant 0$, for a positive integer n such that $n(p+r) = (n+1)(p_0+r)$, the following assertions are equivalent.

(3-1) $A \geqslant B > O$;

(3-12) $B^{\frac{r}{2}}A^{p}B^{\frac{r}{2}} \geqslant (B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^{1+\frac{1}{n}}$;

(3-13) There exists a unique $S > O$ with $\|S\| \leqslant 1$ such that

$$(B^{\frac{r}{2}}A^{p_0}B^{\frac{r}{2}})^{n+1} = ((B^{\frac{r}{2}}A^{p}B^{\frac{r}{2}})^{\frac{1}{2}}S(B^{\frac{r}{2}}A^{p}B^{\frac{r}{2}})^{\frac{1}{2}})^{n} .$$

Proof. Replace A and B by B^{-1} and A^{-1} , respectively, in (3-10) and (3-11) of Theorem 3.3, then (3-12) and (3-13) are obtained.

References

[1] R.G.Douglas, On majorization, factorization, and range inclusion of operators on Hilbert space, Proc. Amer. Math. Soc., 17 (1966), 413-415.

[2] T. Furuta, $A \geqslant B \geqslant O$ assures $(B^r A^p B^r)^{1/q} \geqslant B^{(p+2r)/q}$ for $r \geqslant 0$, $p \geqslant 0$, $q \geqslant 1$ with $(1+2r)q \geqslant p+2r$, Proc. Amer. Math. Soc., 101 (1987), 85-88.

[3] T. Furuta, An extension of the Furuta inequality and Ando-Hiai log majorization, Linear Algebra Appl., 219 (1995), 139-155.

[4] E. Heinz, Beiträge zur Str̈ungstheorie der Spektrallegung, Math. Ann., 123 (1951), 415-438.

[5] C.-S. Lin, On operator order and chaotic operator order for two operators, Linear Algebra Appl., 425 (2007), 1-6.

[6] K. Löwner, Über monotone Matrixfunktionen, Math. Z., 38 (1934), 177-216.

[7] K. Tanahashi, Best possibility of the Furuta inequality, Proc. Amer. Math. Soc., 124 (1996), 141-146.

[8] K. Tanahashi, The best possibility of the grand Furuta inequality, Proc. Amer. Math. Soc., 128 (2000), 511-519.

[9] J. Yuan, Z. Gao, The operator equation $K^p = H^{\frac{\delta}{2}} T^{\frac{1}{2}} (T^{\frac{1}{2}} H^{\delta+r} T^{\frac{1}{2}})^{\frac{p-\delta}{\delta+r}} T^{\frac{1}{2}} H^{\frac{\delta}{2}}$ and its applications, J. Math. Anal. Appl., 341 (2008), 870-875.

[10] J. Yuan, Z. Gao, Classified construction of generalized Furuta type operator functions, Math. Inequal. Appl., 11 (2) (2008), 189-202.

[11] J. Yuan, Z. Gao, Complete form of Furuta inequality, Proc. Amer. Math. Soc., 136 (2008), 2859-2867.

[12] J. Yuan, Furuta inequality and q-hyponormal operators, Oper. Matrices, 4 (3) (2010), 405-415.

[13] J. Yuan, G. Ji, Extensions of Kadison's inequality on positive linear maps, to appear.